

Gauge dependence and renormalization of $\tan \beta$ in the MSSM

Ayres Freitas and Dominik Stöckinger *

*Deutsches Elektronen-Synchrotron DESY,
D-22603 Hamburg, Germany*

Abstract

Well-known and newly developed renormalization schemes for $\tan \beta$ are analyzed in view of three desirable properties: gauge independence, process independence, and numerical stability in perturbation theory. Arguments are provided that no scheme can meet all three requirements, and as an illustration, a “No-Go-Theorem” for the renormalization of $\tan \beta$ is established. Nevertheless, two particularly attractive schemes emerge. A discussion about which scheme might be the best compromise in practice is given.

PACS numbers: 12.60.Jv, 11.10.Gh, 11.15.-q

1 Introduction

The quantity $\tan \beta$ is one of the main input parameters of the minimal supersymmetric standard model (MSSM). At the tree level it is defined as the ratio of the two vacuum expectation values $v_{1,2}$ of the MSSM Higgs doublets,

$$\tan \beta = \frac{v_2}{v_1}. \quad (1)$$

Owing to its central appearance in spontaneous symmetry breaking, $\tan \beta$ plays a crucial role in almost all sectors of the MSSM and has significant impact on most MSSM observables. However, the vacuum expectation values and $\tan \beta$ are not directly measurable quantities. In spite of its importance, $\tan \beta$ is an auxiliary variable. The virtue of $\tan \beta$ is that it can be used as an easy-to-handle input parameter in terms of which all different observables can be expressed. In contrast with other parameters, such as the electron charge or the particle masses, there is no obvious and unique way to relate $\tan \beta$ to an observable.

The actual definition of $\tan \beta$, its physical meaning and its relation to observables is given by the choice of a renormalization scheme. This choice determines the numerical value of $\tan \beta$ as well as its formal properties, such as gauge dependence and renormalization-scale dependence.

*afreitas@mail.desy.de,
dominik@mail.desy.de.

In the literature, several renormalization schemes for $\tan\beta$ have been proposed and used [1–3]. Each of these schemes has specific advantages and disadvantages. In this paper, we give three criteria that are desirable for a renormalization scheme for $\tan\beta$:

- Gauge independence. If the relation between $\tan\beta$ and observables is gauge independent, the numerical value of $\tan\beta$ is also gauge independent and a more physical interpretation of $\tan\beta$ is possible.
- Process independence. If $\tan\beta$ is defined by a relation to a physical process, a non-universality and flavor dependence can be introduced that violates the intuition implied by eq. (1) that $\tan\beta$ is a quantity of the MSSM Higgs sector. Process-dependent schemes have further technical drawbacks like the necessity to calculate more complicated vertex functions in the determination of $\delta\tan\beta$ and the possible appearance of infrared divergent QED or QCD corrections.
- Numerical stability. The numerical properties of the renormalization constant $\delta\tan\beta$ should not spoil the validity of the perturbative expansion. Generally speaking, the finite contribution to $\delta\tan\beta$ and the renormalization-scale dependence of $\tan\beta$ should not be too large.

We analyze the properties of the known renormalization schemes in view of these criteria and devise new schemes in order to satisfy them. Since it is the most intricate point, we will be concerned with questions of gauge dependence and gauge independence for the largest part of the paper.

Our results are negative. We find that the known process-independent schemes are gauge dependent and that the new gauge-independent schemes lead to numerical instabilities. Arguments are provided that schemes satisfying all three criteria do not exist. However, as a result of our analysis two schemes emerge that are particularly attractive and useful compromises.

The outline of the present paper is as follows. After briefly introducing the MSSM Higgs sector in Sec. 2, it is shown in Sec. 3 that the $\overline{DR}$ scheme as well as the schemes proposed in [1, 2] are gauge dependent already at the one-loop level. In Sec. 4, three gauge-independent and process-independent schemes are developed. The numerical instability induced by these schemes is exhibited in Sec. 5. The discussion of process-independent schemes is completed in Sec. 6 by demonstrating that a large class of process- and gauge-independent schemes leads to numerical instabilities. Finally, in Sec. 7 the problems caused by process-dependent schemes are discussed and one useful scheme is presented. Section 8 contains our conclusions and a discussion of two attractive renormalization schemes for $\tan\beta$.

2 The MSSM Higgs sector

The MSSM contains two Higgs doublets

$$H_1 = \begin{pmatrix} v_1 + \frac{1}{\sqrt{2}}(\phi_1 - i\rho_1) \\ -\phi_1^- \end{pmatrix}, \quad H_2 = \begin{pmatrix} \phi_2^+ \\ v_2 + \frac{1}{\sqrt{2}}(\phi_2 + i\rho_2) \end{pmatrix}, \quad (2)$$

whose electrically neutral components are shifted by $v_{1,2}$ in order to account for the finite vacuum expectation values. The parameters of the MSSM Higgs sector are

$$g, g', \tilde{m}_1^2, \tilde{m}_2^2, m_3^2, \quad (3)$$

where g and g' are the SU(2) and U(1) gauge couplings, $\tilde{m}_{1,2}^2 = \mu^2 + m_{1,2}^2$ with the Higgsino parameter μ , and $m_{1,2,3}^2$ are soft-breaking parameters. We assume CP conservation, so all parameters are real. Expressed in terms of these parameters, the Higgs boson potential reads

$$V = \tilde{m}_1^2 |H_1|^2 + \tilde{m}_2^2 |H_2|^2 + m_3^2 (H_1^\dagger H_2^2 - H_1^2 H_2^\dagger + h.c.) + \frac{g^2 + g'^2}{8} (|H_1|^2 - |H_2|^2)^2 + \frac{g^2}{2} |H_1^\dagger H_2|^2. \quad (4)$$

In order to make the physical content more transparent, the parameters of the gauge and Higgs sector (3) and the shifts $v_{1,2}$ are reparametrized in terms of

$$e, M_Z, M_W, M_A, \tan \beta, t_1, t_2 \quad (5)$$

with ($\tilde{g}^2 = g^2 + g'^2$, $v^2 = v_1^2 + v_2^2$)

$$e = \frac{gg'}{\tilde{g}}, \quad M_Z^2 = \frac{\tilde{g}^2 v^2}{2}, \quad M_W^2 = \frac{g^2 v^2}{2}, \quad \tan \beta = \frac{v_2}{v_1}, \quad (6a)$$

$$M_A^2 = \sin^2 \beta \frac{t_1}{\sqrt{2} v_1} + \cos^2 \beta \frac{t_2}{\sqrt{2} v_2} - m_3^2 (\tan \beta + \cot \beta), \quad (6b)$$

$$t_1 = \sqrt{2} \left[\tilde{m}_1^2 v_1 + m_3^2 v_2 + \frac{\tilde{g}^2}{4} v_1 (v_1^2 - v_2^2) \right], \quad (6c)$$

$$t_2 = \sqrt{2} \left[\tilde{m}_2^2 v_2 + m_3^2 v_1 - \frac{\tilde{g}^2}{4} v_2 (v_1^2 - v_2^2) \right]. \quad (6d)$$

For vanishing tadpole parameters $t_1 = t_2 = 0$, M_A^2 is the mass of the pseudoscalar A^0 , and the tree-level Higgs mass eigenstates are given by the fields $H, h, G^0, A^0, G^\pm, H^\pm$ with

$$\begin{pmatrix} \phi_1 \\ \phi_2 \end{pmatrix} = \begin{pmatrix} c_\alpha & -s_\alpha \\ s_\alpha & c_\alpha \end{pmatrix} \begin{pmatrix} H \\ h \end{pmatrix}, \quad (7)$$

$$\begin{pmatrix} \rho_1 \\ \rho_2 \end{pmatrix} = \begin{pmatrix} c_\beta & -s_\beta \\ s_\beta & c_\beta \end{pmatrix} \begin{pmatrix} G^0 \\ A^0 \end{pmatrix}, \quad (8)$$

$$\begin{pmatrix} \phi_1^\pm \\ \phi_2^\pm \end{pmatrix} = \begin{pmatrix} c_\beta & -s_\beta \\ s_\beta & c_\beta \end{pmatrix} \begin{pmatrix} G^\pm \\ H^\pm \end{pmatrix} \quad (9)$$

and

$$\tan 2\alpha = \tan 2\beta \frac{M_A^2 + M_Z^2}{M_A^2 - M_Z^2}, \quad -\pi/2 < \alpha < 0. \quad (10)$$

Here and in the following we use the abbreviations $t_\beta = \tan \beta$, $s_\beta = \sin \beta$, $c_\beta = \cos \beta$, etc.

The tree-level Higgs-boson masses read

$$M_{H^\pm}^{2(0)} = M_A^2 + M_W^2, \quad (11)$$

$$M_{H,h}^{2(0)} = \frac{1}{2} \left(M_A^2 + M_Z^2 \pm \sqrt{(M_A^2 + M_Z^2)^2 - 4M_Z^2 M_A^2 \cos^2(2\beta)} \right). \quad (12)$$

An important equation is given by

$$\cos^2(2\beta) = \frac{M_h^{2(0)} M_H^{2(0)}}{M_A^2 (M_h^{2(0)} + M_H^{2(0)} - M_A^2)}. \quad (13)$$

It relates $\tan \beta$ to the physical Higgs-boson masses and will be useful for later purposes.

The counterterms to the MSSM Higgs sector are generated by a multiplicative renormalization transformation of the parameters and fields. For the purpose of the present paper, the explicit form of the field renormalization is not important, but the parameter renormalization is given by

$$\begin{aligned} g^{(\prime)} &\rightarrow g^{(\prime)} + \delta g^{(\prime)}, & \tilde{m}_{1,2}^2 &\rightarrow \tilde{m}_{1,2}^2 + \delta \tilde{m}_{1,2}^2, \\ m_3^2 &\rightarrow m_3^2 + \delta m_3^2, & v_{1,2} &\rightarrow v_{1,2} + \delta v_{1,2}. \end{aligned} \quad (14)$$

Through the relations (6) also renormalization constants δe , $\delta M_{Z,W,A}$, δt_β and $\delta t_{1,2}$ are defined, in particular $\delta t_\beta = \tan \beta (\frac{\delta \tilde{v}_2}{v_2} - \frac{\delta \tilde{v}_1}{v_1})$. There are two important relations between δt_β and other renormalization constants (for $t_1 = t_2 = 0$) that will be of use later:

$$\frac{\delta t_1}{v_1} + \frac{\delta t_1}{v_2} = \sqrt{2}(\delta \tilde{m}_1^2 + \delta \tilde{m}_2^2) + \sqrt{2}\delta m_3^2(t_\beta + \frac{1}{t_\beta}) + \sqrt{2}m_3^2(1 - \frac{1}{t_\beta^2})\delta t_\beta, \quad (15)$$

$$\delta M_A^2 = -\delta m_3^2(t_\beta + \frac{1}{t_\beta}) - m_3^2(1 - \frac{1}{t_\beta^2})\delta t_\beta + s_\beta^2 \frac{\delta t_1}{\sqrt{2}v_1} + c_\beta^2 \frac{\delta t_2}{\sqrt{2}v_2}. \quad (16)$$

3 Gauge dependence of some schemes for $\tan \beta$

3.1 Definition of the schemes

The actual definition of the parameters (5), their physical meaning and their relations to experimental quantities is given by the choice of a renormalization scheme. In gauge theories a gauge fixing is necessary for quantization. Since the gauge fixing is unphysical, the relations between observable quantities do not depend on it, but the relations between observables and the parameters (5) can be gauge dependent. Accordingly, in such a case the values for the parameters extracted from experiment are gauge dependent.

For e , $M_{Z,W,A}$, the on-shell renormalization scheme (see [4] for the case of the MSSM) provides a gauge-independent definition [5]: e is related to the effective charge in the Thomson limit, $M_{Z,W,A}$ are the masses of the Z , W , A^0 bosons (defined as the real parts of the poles of the respective propagators). In addition, setting the renormalized tadpoles to zero,

$$\Gamma_{\phi_1} = \Gamma_{\phi_2} \stackrel{!}{=} 0, \quad (17)$$

is in agreement with the requirement of gauge independence. Here and henceforth, Γ denotes the generating functional of the renormalized 1PI vertex functions and $\Gamma_{\varphi_1 \dots}(p_1, \dots)$ denotes a vertex function with incoming fields $\varphi_1 \dots$ and incoming momenta $p_1 \dots$.

In contrast, for $\tan \beta$ no such gauge-independent standard-definition is available. In this section we will show explicitly the gauge dependence of several well-known schemes for $\tan \beta$. The first of these is the $\overline{DR}$ scheme, which is defined by the condition

$$\overline{DR} : \quad \delta t_\beta \stackrel{!}{=} \text{pure divergence}, \quad (18)$$

where “pure divergence” denotes a term of the order $\Delta = \frac{2}{4-D} - \gamma_E + \log 4\pi$ in dimensional reduction (or in dimensional regularization when suitable supersymmetry-restoring counterterms have been added [6]) whose prefactor is such that all renormalized quantities are finite. Further common renormalization schemes for $\tan \beta$ are the ones introduced by Dabelstein [1] and by Chankowski et al. [2] (DCPR). In these schemes one writes $\delta \tilde{v}_{1,2} = v_{1,2} \delta Z_{1,2}/2 - \delta v_{1,2}$ with the field renormalization constants $\delta Z_{1,2}$ of the Higgs doublets and requires the conditions

$$\frac{\delta v_1}{v_1} = \frac{\delta v_2}{v_2}, \quad \text{Re} \hat{\Sigma}_{A^0 Z}(M_A^2) = 0, \quad (19)$$

where the renormalized $A^0 Z$ two-point function is decomposed as $\Gamma_{A^0 Z^\mu}(-p, p) = i p_\mu \hat{\Sigma}_{A^0 Z}(p^2)$. At the one-loop level these conditions lead to

$$\text{DCPR} : \quad \delta t_\beta \stackrel{!}{=} \frac{1}{2c_\beta^2 M_Z} \text{Re} \Sigma_{A^0 Z}(M_A^2) \quad (20)$$

with the unrenormalized $A^0 Z$ two-point function $\Sigma_{A^0 Z}$. A similar prescription requiring vanishing $H^+ W^-$ mixing instead of vanishing $A^0 Z$ mixing has been used in [7].

3.2 Extended Slavnov-Taylor identity as a tool

In the presented schemes $\tan \beta$ is apparently not directly related to any observable. It is known that the $\overline{DR}$ scheme leads to a gauge dependent $\tan \beta$ at the two-loop level [8]. In the following, we will show that in fact both schemes lead to a gauge dependence of $\tan \beta$ already at the one-loop level.

The tool we use to determine the gauge dependence is an extended Slavnov-Taylor identity introduced in [9, 10]:

$$\tilde{S}(\Gamma) \equiv S(\Gamma) + \chi \partial_\xi \Gamma = 0. \quad (21)$$

Here ξ denotes an arbitrary gauge parameter in the gauge-fixing term and χ is a fermionic variable acting as the BRS transformation of ξ . $S(\Gamma)$ is the usual Slavnov-Taylor operator (see Appendix). If not stated otherwise we leave the form of the gauge-fixing term open. We only assume that the gauge-fixing term is coupled to auxiliary B fields,

$$\mathbb{L}_{\text{fix}} = \sum_{V=A, Z, W^+, W^-} B^{V\dagger} \mathcal{F}^V + \frac{\xi}{2} |B^V|^2. \quad (22)$$

This simplifies the symmetry identities, but for loop calculations, eq. (22) is equivalent to $\mathbb{L}_{\text{fix}} = -\frac{1}{2\xi} \sum |\mathcal{F}^V|^2$, which is obtained by eliminating the B fields using their equations of motion.

In contrast to the usual Slavnov-Taylor identity $S(\Gamma) = 0$, the extended identity (21) need not be satisfied. If, however, $\tilde{S}(\Gamma) = 0$ holds, then physical quantities — expressed as functions of the parameters of the lowest-order Lagrangian — are gauge independent. Accordingly, $\tilde{S}(\Gamma) = 0$ implies that the values for the parameters extracted from experiment are gauge independent. If suitable χ -dependent terms are added to the action, the extended Slavnov-Taylor identity holds at the tree level. In order to satisfy $\tilde{S}(\Gamma) = 0$ also at higher orders, the renormalization conditions must be chosen so as to not contradict $\tilde{S}(\Gamma) = 0$.

We will make use of the extended Slavnov-Taylor identity in particular in two ways. On the one hand, in our one-loop calculations we employ regularization by dimensional reduction [11] assuming all symmetries are preserved, so $\tilde{S}(\Gamma^{(1),\text{reg}}) = 0$, where Γ^{reg} denotes the unrenormalized vertex functional and the index “(1)” denotes the loop order. The identity $\tilde{S}(\Gamma^{(1),\text{reg}}) = 0$ entails an easy way to calculate the ξ -derivatives $\partial_\xi \Gamma_{\varphi_i \dots}^{(1),\text{reg}}$ of regularized vertex functions.

On the other hand, presuming $\tilde{S}(\Gamma^{\text{reg}}) = 0$ at some loop order, there is a very simple necessary condition for $\tilde{S}(\Gamma) = 0$ at this order, yielding a simple check whether a given set of renormalization conditions is compatible with $\tilde{S}(\Gamma) = 0$. As shown in [10, 12], the counterterms in gauge theories can be generally classified into gauge-dependent total BRS variations and gauge-independent non-BRS variations. Independent of the presence of spontaneous symmetry breaking, the non-BRS variations correspond to the symmetric parameters of the theory, whereas the vacuum expectation values as well as field renormalization constants are related to total BRS variations. This result can be easily transferred to the case of the MSSM Higgs sector (see also [4] for the classification of the MSSM counterterms). It means that in order to satisfy $\tilde{S}(\Gamma) = 0$,

$$\partial_\xi \delta g^{(\prime)} = \partial_\xi \delta \tilde{m}_{1,2}^2 = \partial_\xi \delta m_3^2 = 0 \quad (23)$$

has to hold. In contrast, for $\delta \tilde{v}_{1,2}$ and field renormalization constants a ξ dependence is compatible with $\tilde{S}(\Gamma) = 0$. Equation (23) also constitutes a sufficient condition for $\tilde{S}(\Gamma) = 0$, since if it is satisfied, χ -dependent counterterms can always be added in such a way that $\tilde{S}(\Gamma) = 0$ holds at the considered loop order.

Thus, eq. (23) provides a simple way to check whether a given renormalization scheme is compatible with $\tilde{S}(\Gamma) = 0$. In the following calculations we will use the on-shell conditions for e , $M_{Z,W,A}$ together with the tadpole conditions (17). Since these conditions are compatible with $\tilde{S}(\Gamma) = 0$, eq. (23) provides in particular a check for the renormalization condition for $\tan \beta$ and hence for the gauge independence of $\tan \beta$.

3.3 Calculating the gauge dependence

Let us now show the gauge dependence of $\tan \beta$ at the one-loop level in the $\overline{DR}$ and DCPR schemes (18), (20), beginning with the $\overline{DR}$ scheme. If $\tilde{S}(\Gamma) = 0$ is to hold, the gauge independence of the counterterms (23) implies, together with eq. (15), a certain

gauge dependence of the renormalization constant $\delta t_\beta^{\text{fin}}$:

$$\partial_\xi \left(\frac{\delta t_1}{v_1} + \frac{\delta t_2}{v_2} \right)^{\text{fin}} = \left[\sqrt{2} m_3^2 \left(1 - \frac{1}{t_\beta^2} \right) \right] \partial_\xi \delta t_\beta^{\text{fin}}. \quad (24)$$

We only consider the purely finite parts here since the divergent contributions are restricted by $\tilde{S}(\Gamma^{(1),\text{reg}}) = 0$ and hence are in agreement with $\tilde{S}(\Gamma) = 0$. In the $\overline{DR}$ scheme the finite part of the renormalization constant δt_β itself is zero and therefore gauge independent,

$$\partial_\xi \delta t_\beta^{\text{fin}} = 0, \quad (25)$$

and the question is whether this is compatible with the gauge dependence prescribed by eq. (24).

The tadpole counterterms in (24) are determined by $\Gamma_{\phi_{1,2}} \stackrel{!}{=} 0$ as

$$\delta t_{1,2} = \Gamma_{\phi_{1,2}}^{(1),\text{reg}}. \quad (26)$$

Using the extended Slavnov-Taylor identity at the regularized level, $\tilde{S}(\Gamma^{(1),\text{reg}}) = 0$, yields for the l.h.s. of (24):

$$\begin{aligned} \partial_\xi \left(\frac{\delta t_1}{v_1} + \frac{\delta t_2}{v_2} \right) &= - \sum_{j=1,2} \Gamma_{\chi Y_{\phi_j}}^{(1),\text{reg}} \sum_{i=1,2} \frac{1}{v_i} \Gamma_{\phi_j \phi_i}^{(0)} \\ &= m_3^2 \left(\frac{1}{v_1^2} - \frac{1}{v_2^2} \right) \left(-v_2 \Gamma_{\chi Y_{\phi_1}}^{(1),\text{reg}} + v_1 \Gamma_{\chi Y_{\phi_2}}^{(1),\text{reg}} \right), \end{aligned} \quad (27)$$

where the Y_{ϕ_i} denote the sources of the BRS transformations of ϕ_i used in the Slavnov-Taylor identity. The ξ dependence of $\Gamma_{\phi_i}^{(1),\text{reg}}$ and the results for $\Gamma_{\chi Y_{\phi_i}}^{(1),\text{reg}}$ depend on the specific choice of the gauge fixing. In the R_ξ gauge we obtain (for the relevant Feynman rules see the Appendix):

$$\begin{aligned} \Gamma_{\chi Y_{\phi_i}}^{(1),\text{reg}} &= \begin{pmatrix} \cos \beta \\ \sin \beta \end{pmatrix} \frac{1}{16\pi^2} \left(\frac{\tilde{g} M_Z}{4} B_0(0, \xi M_Z^2, \xi M_Z^2) \right. \\ &\quad \left. + \frac{g M_W}{2} B_0(0, \xi M_W^2, \xi M_W^2) \right), \end{aligned} \quad (28)$$

where B_0 denotes the usual two-point function. Hence, in the context of the R_ξ gauge,

$$\partial_\xi \left(\frac{\delta t_1}{v_1} + \frac{\delta t_2}{v_2} \right)^{\text{fin}} = 0, \quad (29)$$

and the $\overline{DR}$ scheme is compatible with eqs. (24), (25) and hence with $\tilde{S}(\Gamma) = 0$ at the one-loop level. This confirms the result of [8], where one-loop gauge independence, but two-loop gauge dependence of $\tan \beta$ was found in the R_ξ gauge.

However, in more general gauges (29) is not true. As a simple example we consider an infinitesimal deviation from the R_ξ gauge by introducing a second, infinitesimal gauge parameter ζ^{ZA^0} that enters the gauge-fixing function for the Z boson as follows:

$$\mathcal{F}^Z = \partial_\mu Z^\mu + M_Z (\xi G^0 + \zeta^{ZA^0} A^0). \quad (30)$$




Figure 1: The one-loop diagrams to the vertex functions $\Gamma_{\chi Y_{\phi_i}}^{(1),\text{reg}}$ and $\Gamma_{\chi^{ZA^0} Y_{\phi_i}}^{(1),\text{reg}}$.

We can study the dependence on this second gauge parameter in the same way as the dependence on ξ . We introduce the variable χ^{ZA^0} and consider the Slavnov-Taylor identity $\tilde{S}(\Gamma) \equiv S(\Gamma) + \chi^{ZA^0} \partial_{\zeta^{ZA^0}} \Gamma = 0$. As in the general case, the on-shell conditions for e , $M_{Z,W,A}$ and the tadpole conditions (17) are in agreement with $\tilde{S}(\Gamma) = 0$.¹ Using this identity we obtain an identity prescribing the ζ^{ZA^0} dependence of $\delta t_\beta^{\text{fin}}$, analogously to eq. (24),

$$\partial_{\zeta^{ZA^0}} \left(\frac{\delta t_1}{v_1} + \frac{\delta t_2}{v_2} \right)^{\text{fin}} = \left[\sqrt{2} m_3^2 \left(1 - \frac{1}{t_\beta^2} \right) \right] \partial_{\zeta^{ZA^0}} \delta t_\beta^{\text{fin}}. \quad (31)$$

where, analogously to eq. (27), the l.h.s. is given by

$$\partial_{\zeta^{ZA^0}} \left(\frac{\delta t_1}{v_1} + \frac{\delta t_2}{v_2} \right) = m_3^2 \left(\frac{1}{v_1^2} - \frac{1}{v_2^2} \right) \left(-v_2 \Gamma_{\chi^{ZA^0} Y_{\phi_1}}^{(1),\text{reg}} + v_1 \Gamma_{\chi^{ZA^0} Y_{\phi_2}}^{(1),\text{reg}} \right). \quad (32)$$

However, the one-loop results of the vertex functions involving χ^{ZA^0} are (evaluated at $\zeta^{ZA^0} = 0$)

$$\Gamma_{\chi^{ZA^0} Y_{\phi_i}}^{(1),\text{reg}} = \begin{pmatrix} -\sin \beta \\ \cos \beta \end{pmatrix} \frac{1}{16\pi^2} \frac{\tilde{g} M_Z}{2} B_0(0, \xi M_Z^2, M_A^2). \quad (33)$$

Hence, the l.h.s. of eq. (31) is non-vanishing:

$$\partial_{\zeta^{ZA^0}} \left(\frac{\delta t_1}{v_1} + \frac{\delta t_2}{v_2} \right)^{\text{fin}} \neq 0. \quad (34)$$

Thus, in the generalized gauge, the ζ^{ZA^0} independence of $\tan \beta$ and $\tilde{S}(\Gamma) = 0$ imply

$$\partial_{\zeta^{ZA^0}} \delta t_\beta^{\text{fin}} \neq 0. \quad (35)$$

¹Clearly, (30) induces a gauge-dependent mixing of A^0 with G^0 , Z^μ and B^Z . One might wonder whether this mixing has an effect on the on-shell condition for M_A and whether it is still in agreement with the requirement of gauge independence. As shown in [4], as long as the auxiliary B fields are not eliminated from the gauge fixing (22), this on-shell condition can be written as

$$\det \begin{pmatrix} \Gamma_{G^0 G^0} & \Gamma_{G^0 A^0} \\ \Gamma_{A^0 G^0} & \Gamma_{A^0 A^0} \end{pmatrix} (p^2 = \mathcal{M}_A^2) = 0$$

with $\text{Re} \mathcal{M}_A^2 = M_A^2$. Moreover, with uneliminated B fields there is no contribution of the gauge fixing to $\Gamma_{G^0 G^0}$, $\Gamma_{G^0 A^0}$, $\Gamma_{A^0 A^0}$, and the evaluation of the renormalization conditions at the tree level is unchanged compared to the R_ξ case. At higher orders, as shown in [4], this condition for M_A indeed corresponds to the (gauge-independent) pole of the A^0 propagator, even though the A^0 field mixes with B^Z at the tree level — or with G^0 and Z^μ after the elimination of B^Z .




Figure 2: The one-loop diagrams to the vertex functions $\Gamma_{\chi A^0 Y_Z}^{(1),\text{reg}}$ and $\Gamma_{\chi A^0 Y_{G^0}}^{(1),\text{reg}}$.

This is incompatible with the $\overline{DR}$ -condition, which implies $\partial_{\zeta Z A^0} \delta t_\beta^{\text{fin}} = 0$. Conversely, if the $\overline{DR}$ scheme is used, the relation between $\tan \beta$ and observable quantities, i.e. the value $\tan \beta^{\text{exp}}$ extracted from experiment is gauge dependent:

$$\partial_{\zeta Z A^0} \tan \beta_{\overline{DR}}^{\text{exp}} \neq 0. \quad (36)$$

The different result for eq. (33) compared to eq. (28) can be understood by considering the corresponding Feynman diagrams in Fig. 1 (the Feynman rules can be read off from the Lagrangian given in the Appendix). In the R_ξ gauge fixing only the Goldstone bosons $G^{0,\pm}$ appear, and correspondingly only Goldstone bosons contribute as internal scalar lines in the diagrams for $\Gamma_{\chi Y_{\phi_i}}$. In the generalized gauge, however, we study the dependence on the A^0 part of the gauge fixing, and correspondingly only A^0 appears as an internal scalar line in $\Gamma_{\chi Z A^0 Y_{\phi_i}}$. The couplings of $G^{0,\pm}$ and A^0 to $\begin{pmatrix} Y_{\phi_1} \\ Y_{\phi_2} \end{pmatrix}$ are proportional to $\begin{pmatrix} \cos \beta \\ \sin \beta \end{pmatrix}$ and $\begin{pmatrix} -\sin \beta \\ \cos \beta \end{pmatrix}$, respectively, which explains the results (33) and (28).

In a second step it is quite easy to see the gauge dependence of $\tan \beta$ in the DCPR schemes defined by eq. (20). We know that the $\overline{DR}$ scheme is in agreement with $\tilde{S}(\Gamma) = 0$ in the R_ξ gauge at the one-loop level. So we consider the difference

$$\partial_\xi (\delta t_{\beta \text{DCPR}} - \delta t_{\beta \overline{DR}}) \propto \partial_\xi \text{Im} \Gamma_{A^0 Z}^{(1),\text{reg,fin}}(M_A^2). \quad (37)$$

Using $\tilde{S}(\Gamma^{\text{reg}}) = 0$, in the R_ξ gauge the r.h.s. can be expressed as

$$-\left[\sum_{i=1,2} \Gamma_{\chi Y_{\phi_i}}^{(1),\text{reg}} \Gamma_{A^0 Z \phi_i}^{(0)} + \sum_{\varphi=Z, G^0} \Gamma_{\chi A^0 Y_\varphi}^{(1),\text{reg}} \Gamma_{Z \varphi}^{(0)} \right]^{\text{fin}} \neq 0, \quad (38)$$

which is non-zero as can be easily seen by inspection of the corresponding one-loop diagrams (see Fig. 2). For example, there is an M_h -dependent contribution to $\Gamma_{\chi A^0 Y_{G^0}}$ that cannot be cancelled in eq. (38).

Hence, in the DCPR schemes $\tan \beta$ is gauge dependent even at the one-loop level in the R_ξ gauge:

$$\partial_\xi \tan \beta_{\text{DCPR}}^{\text{exp}} \neq 0. \quad (39)$$

4 Three gauge-independent schemes for $\tan \beta$

In the following we present three gauge-independent renormalization schemes for $\tan \beta$. These schemes share the additional property that $\tan \beta$ is defined via quantities in the Higgs sector and without reference to a specific physical process — in that

sense they are closely related to the tree-level definition $\tan \beta = \frac{v_2}{v_1}$. The use of these schemes is on the one hand to demonstrate the existence of such gauge-independent schemes and on the other hand to illustrate three ways to devise gauge-independent renormalization schemes.

The first scheme is defined by the requirement that the parameter m_3^2 and its counterterm is gauge independent as dictated by eq. (23), which is most easily realized by $\overline{DR}$ renormalization:

$$m_3 \text{ scheme :} \quad \delta m_3^2 \stackrel{!}{=} \text{pure divergence.} \quad (40)$$

We refer to this scheme as the “ m_3 scheme”. Together with the on-shell conditions for $e, M_{Z,W,A}, t_{1,2}$ this condition fixes the seventh parameter in (5), $\tan \beta$. Since at the regularized level $\tilde{S}(\Gamma^{(1),\text{reg}}) = 0$ holds, the divergent part of δm_3^2 is gauge independent, hence

$$\partial_\xi \delta m_3^2 = 0 \quad (41)$$

in the m_3 scheme, and thus this scheme is compatible with eq. (23) and $\tilde{S}(\Gamma) = 0$. Hence, it defines $\tan \beta$ in a gauge-independent way. Using the relation (6b) between M_A^2 , $\tan \beta$ and m_3^2 we can derive a result for the finite part of δt_β :

$$\delta M_A^{2\text{fin}} = \sin^2 \beta \frac{\delta t_1^{\text{fin}}}{\sqrt{2}v_1} + \cos^2 \beta \frac{\delta t_2^{\text{fin}}}{\sqrt{2}v_2} - m_3^2(1 - \cot^2 \beta) \delta t_\beta^{\text{fin}}. \quad (42)$$

In this way δt_β is expressed via δM_A^2 and $\delta t_{1,2}$ by the A^0 self energy and the tadpole contributions. The superscript “fin” denotes the purely finite part of the renormalization constants. The result for the purely divergent part of δt_β is the same as in all other schemes.

A second gauge-independent scheme can be read off from eq. (24), which is a necessary consequence of $\tilde{S}(\Gamma) = 0$. The most straightforward way to satisfy eq. (24) is to require

$$\text{Tadpole scheme :} \quad \delta t_\beta^{\text{fin}} \stackrel{!}{=} \frac{1}{\sqrt{2}m_3^2(1 - \frac{1}{t_\beta^2})} \left(\frac{\delta t_1}{v_1} + \frac{\delta t_2}{v_2} \right)^{\text{fin}}. \quad (43)$$

This scheme will be referred to as the “Tadpole scheme”. It can also be obtained in a second, instructive way by temporarily replacing the tadpole conditions $\Gamma_{\phi_{1,2}} = 0$ by the conditions

$$\delta t_{1,2} \stackrel{!}{=} 0. \quad (44)$$

In such a scheme, where the tadpoles are not renormalized, the renormalization constants $\delta \tilde{v}_{1,2}$ are gauge independent if $\tilde{S}(\Gamma) = 0$ holds (since in this case $\delta \tilde{v}_{1,2}$ can be entirely expressed in terms of the gauge-independent constants in eq. (23)). Hence also δt_β is gauge independent, and in combination with eq. (44) the $\overline{DR}$ -condition

$$\delta t_\beta^{\text{fin}} \stackrel{!}{=} 0 \quad (45)$$

is in agreement with $\tilde{S}(\Gamma) = 0$.

The connected Green functions and the physical content of the theory do not change by varying $\delta\tilde{v}_{1,2}$ and accordingly $\delta t_{1,2}$, δt_β , etc. while keeping $\delta g^{(\prime)}$, $\delta\tilde{m}_{1,2}^2$, δm_3^2 fixed. Equation (15) yields a relation between $\delta t_{1,2}$, δt_β and these fixed renormalization constants:

$$\delta t_\beta^{\text{fin}} = \frac{1}{\sqrt{2}m_3^2(1 - \frac{1}{t_\beta^2})} \left(\frac{\delta t_1}{v_1} + \frac{\delta t_2}{v_2} \right)^{\text{fin}} + \mathcal{O}(\delta\tilde{m}_{1,2}^2, \delta m_3^2) \quad (46)$$

Therefore, for physical quantities the conditions $\Gamma_{\phi_{1,2}} = 0$ for the tadpoles and eq. (43) for $\tan\beta$ are equivalent to eqs. (44) and (45).

By construction, the Tadpole scheme (43) combined with the on-shell conditions for $e, M_{Z,W,A}, t_{1,2}$ is in agreement with the extended Slavnov-Taylor identity $\tilde{S}(\Gamma) = 0$. Hence it provides another gauge-independent definition of $\tan\beta$.

In the third scheme, $\tan\beta$ is defined as a combination of physical Higgs-boson masses in agreement with the lowest-order result (13):

$$\text{HiggsMass scheme :} \quad \cos^2(2\beta) \stackrel{!}{=} \frac{M_h^2 M_H^2}{M_A^2 (M_H^2 + M_h^2 - M_A^2)}, \quad (47)$$

where $M_{H,h}^2$ denote the physical masses of H, h (defined as the real parts of the poles of the respective propagators). This scheme is denoted as ‘‘HiggsMass scheme’’. Since the physical masses are gauge-independent quantities, this scheme provides manifestly a gauge-independent definition of $\tan\beta$.

The definition (47) is problematic, because due to higher-order corrections the r.h.s. can get larger than unity. In spite of this possibility, evaluating the requirement (47) at the one-loop level yields a condition for δt_β that always has a solution:

$$0 \stackrel{!}{=} \frac{\hat{\Sigma}_{hh}}{M_h^{2(0)}} + \frac{\hat{\Sigma}_{HH}}{M_H^{2(0)}} - \frac{\hat{\Sigma}_{hh} + \hat{\Sigma}_{HH}}{M_Z^2}, \quad (48)$$

where $\hat{\Sigma}_{hh,HH}$ denote the renormalized hh and HH self energies, evaluated on-shell ($p^2 = M_{h,H}^2$, respectively). This equation is linear in δt_β and can therefore always be solved for δt_β as a function of the unrenormalized self energies, δM_A^2 and $\delta t_{1,2}$.

However, the possibility of the r.h.s. in eq. (47) getting larger than unity already signals that this HiggsMass scheme could cause numerical problems. In the next section, the numerical properties of all three gauge-independent schemes will be discussed.

5 Numerical instability of the gauge-independent schemes

Gauge dependence and gauge independence are important conceptual characteristics of renormalization schemes. But in practice it is also mandatory that a renormalization scheme does not lead to numerical instabilities in the quantum corrections to physical processes. The gauge-dependent schemes presented in section 3 have been successfully used in many practical loop calculations. Only in the calculation of the neutral Higgs-boson masses [13] slight numerical instabilities caused by the scheme of [1]

	$\tan \beta = 3$			$\tan \beta = 50$		
	$m_h^{\max}$	large μ	no mixing	$m_h^{\max}$	large μ	no mixing
$\overline{DR}$	-0.06	-0.06	-0.06	-0.17	-0.17	-0.17
m_3	0.81	-0.46	-0.04	285.29	127.11	4.92
Tadpole	4.50	-0.21	1.24	370.73	140.11	34.53

Table 1: $\partial \tan \beta / \partial \log \bar{\mu}$ in the $\overline{DR}$ - m_3 , and Tadpole scheme for various parameter scenarios. We have chosen $M_A = 500\text{GeV}$, and the remaining parameter values are chosen according to [14].

were reported that are avoided using the $\overline{DR}$ scheme. In this section we will demonstrate that the three gauge-independent schemes presented in the foregoing section lead to much more severe numerical problems at the one-loop level.

A straightforward way to analyze the numerical behavior of the one-loop corrections is to derive the renormalization-scale dependence of $\tan \beta$ in the various schemes. In the DCPR schemes (20) and in the HiggsMass scheme (47), $\tan \beta$ is scale independent because it is defined on-shell, while in the $\overline{DR}$ scheme, the m_3 scheme and the Tadpole scheme the scale-dependence of $\tan \beta$ is obtained by simply equating

$$(\tan \beta + \delta t_\beta(\bar{\mu}))_{\text{DCPR}} = (\tan \beta(\bar{\mu}) + \delta t_\beta(\bar{\mu}))_{\text{other scheme}}, \quad (49)$$

where $\bar{\mu}$ is the renormalization scale used in dimensional reduction. For the scale-dependence in the $\overline{DR}$ scheme we obtain a simple analytical formula, which is well-known (see e.g. [8]):

$$\frac{\partial}{\partial \log \bar{\mu}} (\tan \beta(\bar{\mu}))_{\overline{DR}} = \tan \beta \frac{1}{16\pi^2} (3h_b^2 - 3h_t^2 + h_\tau^2), \quad (50)$$

where $h_{t,b,\tau}$ are the Yukawa couplings of the top, bottom, and τ , respectively, and where the contributions of the first two generations are neglected. The analytical results for the scale dependence in the other schemes are more complicated, so we restrict ourselves to a numerical analysis. Table 1 contains the numerical results for the scale dependence for three typical sets of MSSM-parameters taken from [14].

Apparently, the scale dependence of $\tan \beta$ in the $\overline{DR}$ scheme is quite small, indicating a small uncertainty in one-loop corrections due to δt_β . On the other hand, the scale dependence of $\tan \beta$ in the m_3 and Tadpole schemes can get immensely large. In practice, for instance changing the scale $\bar{\mu}$ from m_t to M_A can cause unacceptably large changes in the numerical values of one-loop corrections to observables that depend on $\tan \beta$.

The fact that $\tan \beta$ is not scale-dependent in the HiggsMass scheme does not imply that this scheme does not lead to numerical instabilities. In table 2 the numerical values of the purely finite part $\delta t_\beta^{\text{fin}}$ are shown in the schemes where it is non-vanishing. Typically, the finite contribution to a renormalization constant should be suppressed compared to the respective tree-level parameter. Table 2 shows that this is the case only for the DCPR schemes (and of course for the $\overline{DR}$ scheme), whereas in the three gauge-independent schemes, in particular in the HiggsMass scheme, $\delta t_\beta^{\text{fin}}$ can exceed $\tan \beta$

	$\tan \beta = 3$			$\tan \beta = 50$		
	$m_h^{\max}$	large μ	no mixing	$m_h^{\max}$	large μ	no mixing
DCPR	-0.10	-0.06	-0.08	3.56	14.47	0.46
m_3	0.56	-0.08	-0.04	490.45	-67.14	-4.85
Tadpole	2.64	-0.46	0.33	624.70	-76.46	0.92
HiggsMass	-2.44	-1.83	-1.33	-426.54	-1995.93	-314.50

Table 2: $\delta t_\beta^{\text{fin}}$ in the DCPR, m_3 , Tadpole, and HiggsMass scheme for various parameter scenarios. We have chosen $\bar{\mu} = M_A = 500\text{GeV}$, and the remaining parameter values are chosen according to [14].

by far. The consequences are very large one-loop corrections to quantities depending on δt_β , signaling the breakdown of the validity of the perturbative expansion.

As an example for the influence on the calculation of observables we consider the one-loop results for the mass of the light CP-even Higgs boson, M_h , according to the strict one-loop formula

$$M_h^2 = M_h^{2(0)} - \text{Re} \hat{\Sigma}_{hh}(M_h^{2(0)}), \quad (51)$$

where $\hat{\Sigma}_{hh}$ is the renormalized one-loop h -self energy. The numerical results are presented in table 3 for the case of small $\tan \beta$, where the Higgs-boson mass is most sensitive to $\tan \beta$.

We observe that for the parameters used here the DCPR schemes and the $\overline{DR}$ scheme agree well numerically, whereas the discrepancy between these schemes and the gauge-independent ones can be very large. These discrepancies cannot be interpreted as theoretical errors in the prediction of the Higgs-boson mass but they are consequences of the invalidity of the perturbative expansion in the m_3 , Tadpole and HiggsMass scheme.

In contrast, perturbation theory is trustworthy in the DCPR schemes and the $\overline{DR}$ scheme. A more detailed comparison between the scheme of [1] and the $\overline{DR}$ scheme taking into account the leading two-loop effects has shown that the $\overline{DR}$ scheme has generally a better numerical behavior in certain regions of the parameter space [13].

6 Impossibility of defining $\tan \beta$ in the Higgs sector without introducing gauge dependence or numerical instability

In the previous section severe numerical problems were found in all three gauge-independent schemes. In this section it is shown that the conflict between gauge independence and numerical stability is unavoidable if $\tan \beta$ is defined via quantities of the Higgs sector. Since the numerical instability is so strong that the schemes cannot be used in practice, this result can be interpreted as a “No-Go”-like theorem for a gauge-independent renormalization of $\tan \beta$ in the Higgs sector.

More precisely, we study the class of renormalization schemes where $\delta t_\beta^{\text{fin}}$ at the one-loop level is given by a linear combination of finite parts of the on-shell Higgs self

	$m_h^{\max}$	large μ	no mixing
Tree level	72.51	72.51	72.51
DCPR	134.44	97.40	112.23
$\overline{DR}, \bar{\mu} = M_A$	135.03	97.93	112.83
$\overline{DR}, \bar{\mu} = m_t$	134.63	97.38	112.35
HiggsMass	119.58	81.48	102.87
$m_3, \bar{\mu} = M_A$	138.34	97.28	112.56
$m_3, \bar{\mu} = m_t$	143.23	93.23	112.25
Tadpole, $\bar{\mu} = M_A$	149.94	94.08	115.18
Tadpole, $\bar{\mu} = m_t$	173.45	92.21	123.97

Table 3: The light Higgs mass M_h in the five renormalization schemes and for various parameter scenarios. We have chosen $\tan \beta = 3$ and $M_A = 500\text{GeV}$, and the remaining parameter values are chosen according to [14].

energies, their momentum derivatives Σ' , and tadpoles:

$$\delta t_\beta^{\text{fin}} = \text{linear combination of } \left(\Sigma_{A^0 A^0}(M_A^2), \Sigma_{A^0 G^0}(M_A^2), \Sigma_{A^0 Z}(M_A^2), \Sigma_{HH}(M_H^2), \right. \\ \left. \Sigma_{hh}(M_h^2), \Sigma_{Hh}(M_{H,h}^2), \Sigma'_{A^0 A^0}(M_A^2), \Sigma'_{HH}(M_H^2), \Sigma'_{hh}(M_h^2), \delta t_1, \delta t_2 \right)^{\text{fin}} \quad (52)$$

The coefficients in this linear combination should be functions of the parameters of the Higgs potential $e, \tan \beta, M_{Z,W,A}$. The choice of (52) is motivated by the intuition that $\tan \beta$ is a quantity of the MSSM Higgs sector. All schemes considered in sections 3, 4 belong to this class.

As shown in Appendix C, the most general gauge-independent schemes of the class (52) are given by

$$\delta t_\beta^{\text{fin}} = \delta t_{\beta m_3}^{\text{fin}} + \left(a_A K_A + a_H K_H + a_h K_h \right)^{\text{fin}} \quad (53)$$

where $a_{A,H,h}$ are coefficients and the quantities $K_{A,H,h}$ are defined as the following combinations:

$$K_A = \Sigma_{A^0 A^0} - \sum_{i,j=1,2} \Gamma_{A^0 A^0 \phi_i}^{(0)} \left(\Gamma_{\phi\phi}^{(0)} \right)^{-1}_{ij} \delta t_j \quad (54)$$

and analogous for $K_{H,h}$. These combinations are gauge independent as can be seen from the identities (displayed here for the case of K_A)

$$\partial_\xi \delta t_i = -\Gamma_{\chi Y \phi_j}^{(1),\text{reg}} \Gamma_{\phi_j \phi_i}^{(0)}, \quad \partial_\xi \Sigma_{A^0 A^0} = -\Gamma_{\chi Y \phi_j}^{(1),\text{reg}} \Gamma_{A^0 A^0 \phi_j}^{(0)}, \quad (55)$$

derived from $\tilde{S}(\Gamma^{(1),\text{reg}}) = 0$. Besides, $K_{A,H,h}$ are nothing but the gauge-independent $\delta M_{A,H,h}^2$ mass counterterms in the scheme with $\delta t_{1,2} = 0$ (compare discussion of the Tadpole scheme).

The numerical instability originates from several terms contributing to δt_β and the scale dependence $\partial_{\log \bar{\mu}} \tan \beta$. In particular, in the general scheme (53), $\partial_{\log \bar{\mu}} \tan \beta$ contains terms of the order

$$\frac{\mu^2}{M_A^2}, \quad \frac{M_{\text{Susy}}^2}{M_A^2}, \quad \frac{M_2^2}{M_A^2}, \quad (56)$$

which can get large independently of each other.² It is possible to choose the coefficients $a_{A,H,h}$ in eq. (53) such that these three large terms are exactly cancelled in $\partial_{\log \bar{\mu}} \tan \beta$. In this way an optimal, gauge-independent scheme with minimal scale dependence is obtained. Of course, this optimal scheme is nothing but the HiggsMass scheme (47), where $\tan \beta$ is defined on-shell and therefore $\bar{\mu}$ -independent. As seen in section 5, the HiggsMass scheme leads to numerically not acceptable, large contributions to δt_β itself.

Hence, the only scheme where the contributions (56) are absent is the numerically unacceptable HiggsMass scheme, and all other schemes of the form (53) involve the large contributions (56) to $\partial_{\log \bar{\mu}} \tan \beta$ and are for this reason unsuited. None of the gauge-independent schemes defined in eq. (53) can be used in practice.

7 Process-dependent schemes

If $\tan \beta$ is defined via quantities of the Higgs sector there is an unavoidable conflict between gauge independence and numerical stability. In order to circumvent these problems one could try to define $\tan \beta$ outside of the Higgs sector by relating it to a specific physical process. This method has been adopted in [3], where it was suggested to use the decay $H^+ \rightarrow \tau^+ \nu_\tau$. The one-loop corrected decay width to this process reads

$$\Gamma[H^+ \rightarrow \tau^+ \nu_\tau] = \frac{\alpha m_\tau^2 M_{H^\pm}^2 t_\beta^2}{8 M_W^2 s_W^2} \left[1 + F_{H^+ \tau \nu} + 2 \frac{\delta e}{e} + 2 \frac{\delta m_\tau}{m_\tau} + 2 \frac{\delta t_\beta}{t_\beta} - \frac{\delta M_W^2}{M_W^2} - 2 \frac{\delta s_W}{s_W} \right], \quad (57)$$

where $F_{H^+ \tau \nu}$ is the form factor describing the vertex and external wave-function corrections to the amplitude $H^+ \rightarrow \tau^+ \nu_\tau$ including the $H^+ - W^+$ and $H^+ - G^+$ mixing self energies.

By requiring that the radiatively corrected decay width retains the same form as the lowest-order formula, eq. (57) can be understood as a definition of the renormalization constant δt_β . As a consequence of the relation to a physical observable, this definition of $\tan \beta$ is manifestly gauge independent.

However, this scheme also has several drawbacks. At first, for the computation of δt_β it is necessary to compute loop corrections to the three-particle vertex in $F_{H^+ \tau \nu}$, which can be difficult beyond the one-loop level. Furthermore, it is conceptually disadvantageous to define $\tan \beta$ in a specific process, since in this way it becomes a non-universal, flavor-dependent quantity.

²In many of the cases in tab. 1, 2, 3 these terms are actually subdominant due to the large value of M_A . Nevertheless, in general they are very significant, especially if the ratios in eq. (56) get large.

Finally, the decay vertex $H^+ \rightarrow \tau^+ \nu_\tau$ also receives QED corrections, which necessarily include contributions with real photon emission in order to cancel infrared divergences. It is not possible to separate the QED corrections from the rest of the electroweak corrections since they are not individually UV finite. For practical calculations, however, it is unacceptable to include real bremsstrahlung corrections into the definition of a counterterm, since this procedure should depend on experimental phase-space cuts and is therefore technically very involved.

While the first two drawbacks hold for any process-dependent scheme, the problem posed by the QED corrections can be avoided by the choice of another process. One possibility is given by the decay $A^0 \rightarrow \tau^+ \tau^-$. Its one-loop decay width reads

$$\Gamma[A^0 \rightarrow \tau^+ \tau^-] = \frac{\alpha m_\tau^2 M_A^2 t_\beta^2}{8 M_W^2 s_W^2} \left[1 + F_{A^0 \tau \tau} + 2 \frac{\delta e}{e} + 2 \frac{\delta m_\tau}{m_\tau} + 2 \frac{\delta t_\beta}{t_\beta} - \frac{\delta M_W^2}{M_W^2} - 2 \frac{\delta s_W}{s_W} \right]. \quad (58)$$

The QED corrections to this decay width consist of the photon loop contributions to $F_{A^0 \tau \tau}$ and δm_τ ,

$$F_{A^0 \tau \tau} = F_{A^0 \tau \tau}^{\text{QED}} + F_{A^0 \tau \tau}^{\text{weak}}, \quad \delta m_\tau = \delta m_\tau^{\text{QED}} + \delta m_\tau^{\text{weak}}. \quad (59)$$

These QED corrections $F_{A^0 \tau \tau}^{\text{QED}} + 2 \frac{\delta m_\tau^{\text{QED}}}{m_\tau}$ form a UV-finite subset of the full electroweak one-loop corrections. The reason for this difference to the decay $H^+ \rightarrow \tau^+ \nu_\tau$ is that the latter process relies substantially on the SU(2) symmetry and γ , Z and W loops have to be summed to yield a UV-finite result. In contrast, the QED corrections to $A^0 \rightarrow \tau^+ \tau^-$ can be thought of as being generated by an effective theory containing essentially the $A^0 \tau \tau$ vertex and the photon and are therefore naturally finite.

Owing to the finite QED corrections, a possible definition of $\tan \beta$ is given by requiring that the pure weak corrections in eq. (58) cancel and that the exact decay width is given by the tree-level result plus QED corrections:

$$\left[1 + F_{A^0 \tau \tau}^{\text{weak}} + 2 \frac{\delta e}{e} + 2 \frac{\delta m_\tau^{\text{weak}}}{m_\tau} + 2 \frac{\delta t_\beta}{t_\beta} - \frac{\delta M_W^2}{M_W^2} - 2 \frac{\delta s_W}{s_W} \right] \stackrel{!}{=} 1, \quad (60)$$

$$\Gamma[A^0 \rightarrow \tau^+ \tau^-] = \frac{\alpha m_\tau^2 M_A^2 t_\beta^2}{8 M_W^2 s_W^2} \left[1 + F_{A^0 \tau \tau}^{\text{QED}} + 2 \frac{\delta m_\tau^{\text{QED}}}{m_\tau} \right]. \quad (61)$$

With this definition, $\tan \beta$ is gauge independent and not affected by QED-corrections and infrared divergences. Moreover, the one-loop correction to the decay $A^0 \rightarrow \tau^+ \tau^-$ using the renormalization scheme of [1] is quite small [15]. Hence, if $\tan \beta$ is defined by eq. (60), it does not suffer from the numerical instabilities found for the cases of the gauge-independent schemes in sections 5, 6.

8 Conclusions

In this paper, the renormalization of $\tan \beta$ has been studied in view of the criteria gauge independence, process independence, and numerical stability. The $\overline{DR}$ scheme as well as the schemes presented in [1, 2] have been shown to imply a gauge dependence of

$\tan \beta$ already at the one-loop level. Therefore, three gauge-independent schemes have been developed — however, using these schemes produces unacceptably large numerical instabilities in higher-order calculations. The conflict between gauge independence and numerical stability has been made more explicit by a general statement about a large class of process-independent schemes, where $\tan \beta$ is defined via quantities of the Higgs sector. We have shown that all gauge-independent schemes of this class lead to numerical instabilities. Hence, it seems to be impossible to find any renormalization prescription that satisfies all three above criteria.

As a way out of these problems, process-dependent schemes can be used, but such schemes also have specific drawbacks. Conceptually, the flavor dependence of $\tan \beta$ is unsatisfactory, and technically, the necessity to calculate three-point functions and the possible appearance of QED and QCD corrections are disadvantages.

In the course of our analysis, two schemes emerge as the best compromises both in conceptual and practical respects. On the one hand, the $\overline{DR}$ scheme is the most advantageous among the process-independent schemes. It is technically very convenient, numerically perfectly well behaved — and although it is in general gauge dependent, it is not gauge dependent at the one-loop level in the context of the important class of R_ξ gauges. On the other hand, defining $\tan \beta$ via the decay $A^0 \rightarrow \tau^+ \tau^-$ provides a particularly attractive process-dependent alternative. In this scheme, $\tan \beta$ is directly connected to an observable and therefore gauge independent as well as renormalization-scale independent. Furthermore, this specific process is theoretically very clean since it involves no QCD corrections at the one-loop level and the QED corrections can be split off.

Both the $\overline{DR}$ scheme and the $(A^0 \rightarrow \tau^+ \tau^-)$ scheme have specific advantages, so depending on the situation one or the other can be more useful in practice. However, finally a decision should be made for one definition of $\tan \beta$, since a common renormalization scheme is important to allow direct comparisons between different higher-order calculations. In the $(A^0 \rightarrow \tau^+ \tau^-)$ scheme, the specific process is chosen merely for technical reasons. From an experimental point of view, the decay $A^0 \rightarrow \tau^+ \tau^-$ is only one possibility amongst a variety of potential observables for the determination of $\tan \beta$ (see e.g. [16]); a key observable for the definition of $\tan \beta$ does not exist. This reflects the fact that $\tan \beta$ is an auxiliary parameter. Accordingly, the advantages of the process-dependent scheme appear less significant. Owing to its technical convenience and its process independence, we assess the $\overline{DR}$ scheme as the best choice for defining $\tan \beta$.

Acknowledgments: We are grateful to J. Guasch and W. Hollik for valuable discussions.

A Slavnov-Taylor operator

The (gauge part of the) Slavnov-Taylor operator of the MSSM can be written in the form

$$S(\Gamma) = \int d^4x \left[\sum_{\substack{\varphi=\phi_{1,2}, G^0, A^0, G^\pm, H^\pm, \\ \text{other fields}}} \frac{\delta \Gamma}{\delta Y_\varphi} \frac{\delta \Gamma}{\delta \varphi} \right]$$

$$+ \sum_{V=A,Z,W^+,W^-} \left(\frac{\delta\Gamma}{\delta Y_V^\mu} \frac{\delta\Gamma}{\delta V_\mu} + \frac{\delta\Gamma}{\delta Y_{c^V}} \frac{\delta\Gamma}{\delta c^V} + B^V \frac{\delta\Gamma}{\delta \bar{c}^V} \right) \Big], \quad (62)$$

where “other fields” stands for the (s)quark, (s)lepton, chargino and neutralino fields. For the full form also including supersymmetry and translational ghosts, see ref. [4]. The fields B^V are auxiliary fields that couple to the gauge-fixing term and are defined as the BRS transformation of the Faddeev-Popov antighosts $\bar{c}^V$. The fields Y_{φ_i} are sources for the BRS transformations of the fields φ_i . For our purposes, the Feynman rules involving $Y_{Z,G^0,A^0,\phi_{1,2}}$ are important. They can be derived from the Lagrangian (s denotes the generator of BRS transformations):

$$\begin{aligned} \mathbb{L}_Y &= Y_{\phi_1} s\phi_1 + Y_{\phi_2} s\phi_2 + Y_{G^0} sG^0 + Y_{A^0} sA^0 + Y_{Z^\mu} sZ^\mu + \dots \\ &= Y_{\phi_1} \frac{1}{2} [g (ic^+(c_\beta G^- - s_\beta H^-) + h.c.) + \tilde{g} c^Z (-c_\beta G^0 + s_\beta A^0)] \\ &\quad + Y_{\phi_2} \frac{1}{2} [g (-ic^+(s_\beta G^+ + c_\beta H^+) + h.c.) + \tilde{g} c^Z (-s_\beta G^0 - c_\beta A^0)] \\ &\quad + Y_{G^0} \frac{1}{2} [g (c^+(-G^-) + h.c.) + \tilde{g} c^Z (c_\beta \phi_1 + s_\beta \phi_2) + 2c^Z M_Z] \\ &\quad + Y_{A^0} \frac{1}{2} [g (c^+(-H^-) + h.c.) + \tilde{g} c^Z (-s_\beta \phi_1 + c_\beta \phi_2)] \\ &\quad + Y_{Z^\mu} [\partial^\mu c^Z - g_{cW} (-iW^+ c^- + iW^- c^+)] + \dots \end{aligned} \quad (63)$$

with $\tilde{g} = \sqrt{g^2 + g'^2}$ and $c_W = \frac{M_W}{M_Z}$.

B Gauge-fixing terms

For most parts of the paper we leave the form of the gauge fixing open. However, in the one-loop calculations two specific gauge-fixing terms are used. The R_ξ gauge is defined by the choice

$$\mathcal{F}^A = \partial^\mu A_\mu, \quad (64)$$

$$\mathcal{F}^Z = \partial^\mu Z_\mu + M_Z \xi G^0, \quad (65)$$

$$\mathcal{F}^\pm = \partial^\mu W_\mu^\pm \pm i M_W \xi G^\pm, \quad (66)$$

$$\mathbb{L}_{\text{fix}} = \sum_{V=A,Z,W^+,W^-} B^{V\dagger} \mathcal{F}^V + \frac{\xi}{2} |B^V|^2 \quad (67)$$

for the gauge-fixing functions and the gauge-fixing Lagrangian. For loop calculations, it is useful to eliminate the B fields via their equations of motion, yielding $\mathbb{L}_{\text{fix}} = -\frac{1}{2\xi} \sum |\mathcal{F}^V|^2$. The χ -dependent terms (as well as the ghost terms) in the Lagrangian that are necessary to satisfy $\tilde{S}(\Gamma^{(0)}) = 0$ are obtained from $\mathbb{L}_{\text{fix,ghost},\chi} = (s + \chi \partial_\xi)(\bar{c}^V (\mathcal{F}^V + \frac{\xi}{2} B^V))$, where s denotes the BRS operator. For eliminated B fields they read

$$\begin{aligned} \mathbb{L}_\chi^{R_\xi\text{-gauge}} &= -\frac{1}{2\xi} \chi \left[\bar{c}^A \partial^\mu A_\mu + \bar{c}^Z (\partial^\mu Z_\mu - \xi M_Z G^0) \right. \\ &\quad \left. + \bar{c}^- (\partial^\mu W_\mu^+ - i\xi M_W G^+) + \bar{c}^+ (\partial^\mu W_\mu^- + i\xi M_W G^-) \right]. \end{aligned} \quad (68)$$

In the generalized gauge defined by eq. (30), the correct form of the χ^{ZA^0} -dependent terms is given by

$$\mathbb{L}_{\chi^{ZA^0}} = \chi^{ZA^0} \bar{c}^Z M_Z A^0. \quad (69)$$

C General gauge-independent scheme

In this section we show that the most general gauge-independent renormalization scheme of the class (52) is given by eq. (53).

Since the m_3 scheme is gauge independent, any other gauge-independent scheme has to satisfy

$$\partial_\xi(\delta t_\beta - (\delta t_\beta)_{m_3}) = 0, \quad (70)$$

leading us to the question on which combinations of the quantities in eq. (52) are gauge independent. In order to answer this question it is sufficient to study linear combinations of the quantities in eq. (52), leaving away $\Sigma_{A^0 A^0, HH, hh}$ and $\Sigma_{A^0 Z}$ since the gauge-independent combinations $K_{A, H, h}$ involving $\Sigma_{A^0 A^0, HH, hh}$ are already known and since $\Sigma_{A^0 Z}$ itself is a linear combination of $\Sigma_{A^0 G^0}$ and $\delta t_{1,2}$ as a consequence of the Slavnov-Taylor identity.

The gauge dependence of the remaining quantities can be expressed using the identity $\tilde{S}(\Gamma^{(1), \text{reg}}) = 0$ as

$$\begin{aligned} & \partial_\xi (\delta t_1, \delta t_2, \Sigma_{A^0 G^0}, \Sigma_{Hh}(M_H^2), \Sigma_{Hh}(M_h^2), \Sigma'_{A^0 A^0}, \Sigma'_{HH}, \Sigma'_{hh})^T \\ &= \mathcal{M} \left(\Gamma_{\chi Y_{\phi_1}}^{(1), \text{reg}}, \Gamma_{\chi Y_{\phi_2}}^{(1), \text{reg}}, \Gamma_{\chi A^0 Y_{G^0}}^{(1), \text{reg}}, \Gamma_{G^0 G^0}^{(0)} + \Gamma_{\chi A^0 Y_{Z^\mu}}^{(1), \text{reg}}, \Gamma_{G^0 Z^\mu}^{(0)}, \right. \\ & \quad \left. \Gamma_{\chi H Y_h}^{(1), \text{reg}}(M_H^2), \Gamma_{\chi h Y_H}^{(1), \text{reg}}(M_h^2), \Gamma_{\chi A^0 Y_{A^0}}^{(1), \text{reg}}, \Gamma_{\chi H Y_H}^{(1), \text{reg}}, \Gamma_{\chi h Y_h}^{(1), \text{reg}} \right)^T, \quad (71) \end{aligned}$$

where $\mathcal{M}$ denotes an invertible matrix whose entries consist of tree-level expressions like $\Gamma_{\phi_i \phi_j}^{(0)}$ etc.

Calculation of the vertex functions involving χ shows that in the R_ξ gauge there is only one vanishing linear combination, namely $s_\beta \Gamma_{\chi Y_{\phi_1}}^{(1), \text{reg}} - c_\beta \Gamma_{\chi Y_{\phi_2}}^{(1), \text{reg}} = 0$. The other vertex functions involving χ in eq. (71) are linearly independent. Correspondingly, the combination (29) is the only gauge-independent linear combination of the quantities on the l.h.s. of eq. (71) in the R_ξ gauge. However, in more general gauges this combination is not gauge independent (see eq. (34)). Hence, there is no gauge-independent linear combination of the quantities on the l.h.s. of eq. (71).

This confirms that the expressions $K_{A, H, h}$ are the only gauge-independent combinations of the quantities in eq. (52), as was to be shown.

References

- [1] A. Dabelstein, *Z. Phys.* **C67** (1995) 495.
- [2] P. Chankowski, S. Pokorski, J. Rosiek, *Nucl. Phys.* **B423** (1994) 437.

- [3] J.A. Coarasa, D. Garcia, J. Guasch, R.A. Jiménez, J. Solà, *Eur. Phys. J.* **C2** (1998) 373.
- [4] W. Hollik, E. Kraus, M. Roth, C. Rupp, K. Sibold, D. Stöckinger, *Nucl. Phys.* **B639** (2002) 3.
- [5] P. Gambino, P.A. Grassi, *Phys. Rev.* **D62** (2000) 076002.
- [6] S.P. Martin, M.T. Vaughn, *Phys. Lett.* **B318** (1993) 331; W. Beenakker, R. Höpker, P.M. Zerwas, *Phys. Lett.* **B378** (1996) 159; W. Hollik, E. Kraus, D. Stöckinger, *Eur. Phys. J.* **C11** (1999) 365; W. Hollik, D. Stöckinger, *Eur. Phys. J.* **C20** (2001) 105.
- [7] L.H. Wan, W.G. Ma, R.Y. Zhang and J. Yi, *Phys. Rev.* **D64** (2001) 115004.
- [8] Y. Yamada, *Phys. Lett.* **B530** (2002) 174.
- [9] H. Kluberg-Stern, J.B. Zuber, *Phys. Rev.* **D12** (1975) 467,482,3159.
- [10] O. Piguet, K. Sibold, *Nucl. Phys.* **B253** (1985) 517.
- [11] W. Siegel, *Phys. Lett.* **B84** (1979) 193.
- [12] R. Häussling, E. Kraus, *Z. Phys.* **C75** (1997) 739.
- [13] M. Frank, S. Heinemeyer, W. Hollik, G. Weiglein, **hep-ph/0202166**.
- [14] M. Carena, S. Heinemeyer, C. Wagner, G. Weiglein, **hep-ph/9912223**.
- [15] A. Dabelstein, *Nucl. Phys.* **B456** (1995) 25.
- [16] J.F. Gunion, T. Han, J. Jiang, S. Mrenna, A. Sopczak, in *Proc. of the APS/DPF/DPB Summer Study on the Future of Particle Physics (Snowmass 2001)* ed. R. Davidson and C. Quigg, **hep-ph/0112334**.